



Note

Structure of 4-connected claw-free graphs not containing a subdivision of K_5 Roi Krakovski^{a,*}, D. Christopher Stephens^b^a Department of Mathematics, Simon Fraser University, Burnaby, B.C. V5A 1S6, Canada^b Department of Mathematical Sciences, Middle Tennessee State University, Murfreesboro, TN 37132, USA

ARTICLE INFO

Article history:

Received 23 May 2010

Received in revised form 14 February 2011

Accepted 15 February 2011

Available online 22 March 2011

Keywords:

Subdivisions of K_5

Claw-free graphs

4-connected

ABSTRACT

We characterize all the 4-connected claw-free graphs not containing a subdivision of K_5 . We show that such graphs are either planar or can be constructed in a simple way starting from a triangle.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

All graphs in this paper are finite and simple. Paths and cycles are simple (that is, have no “repeated vertices”) and are considered subgraphs.

For $X \subseteq V(G)$, the subgraph $G[X]$ induced on X is the subgraph with vertex set X and edge set all edges of G with both ends in X . ($V(G)$ and $E(G)$ denote the vertex set and edge set of G respectively.) We say that $X \subseteq V(G)$ is a *claw* in G if $|X| = 4$ and $G[X]$ is isomorphic to the complete bipartite graph $K_{1,3}$. We say G is claw-free if no $X \subseteq V(G)$ is a claw in G .

If G is a graph and $A \subseteq G$, then $N_G(A)$ (or simply $N(A)$) denotes the set of vertices in $V(G) - V(A)$ adjacent to some vertex in $V(A)$. If $A = \{a\} \subseteq V(G)$, then we use $N_G(a)$ or $N(a)$ instead of $N_G(\{a\})$ or $N(\{a\})$.

If a path P has end vertices u and v , then P is called a u, v -path. For a graph G , and $A_i \subseteq V(G)$ or $A_i \subseteq G$ for $i = 1, 2$, we say P is an A_1, A_2 -path if P is an a_1, a_2 -path with $V(P) \cap V(A_i) = \{a_i\}$ for $i = 1, 2$. We say P is an A_1 -path if P is an a_1, a_2 -path with $V(P) \cap V(A_1) = \{a_1, a_2\}$.

We say that $S \subseteq V(G)$ *separates* A_1 from A_2 if every A_1, A_2 -path contains a vertex from S . A *separation* of a graph G is a pair of subgraphs (G_1, G_2) of G , such that $E(G_1) \cap E(G_2) = \emptyset$, $G_1 \cup G_2 = G$, and $|V(G_1)|, |V(G_2)| > |V(G_1) \cap V(G_2)|$. The order of a separation (G_1, G_2) is $|V(G_1) \cap V(G_2)|$. A k -separation is a separation of order k . Observe that a graph G is k -connected if $|V(G)| > k$ and G has no k' -separation for any $k' < k$.

Let $k \geq 3$ be an integer and let $k_1 = \lceil k/2 \rceil$ and $k_2 = \lfloor k/2 \rfloor$. We say that a graph H is a k -triangular-chain if $V(H) = \{v_1, \dots, v_{k_1}, u_1, \dots, u_{k_2}\}$ and $E(H) = A_1 \cup A_2 \cup A_3 \cup A_4$, where $A_1 = \bigcup_{i=1}^{k_1-1} v_i v_{i+1}$, $A_2 = \bigcup_{i=1}^{k_2-1} u_i u_{i+1}$, $A_3 = \bigcup_{i=1}^{k_2} v_i u_i$, $A_4 = \bigcup_{i=1}^{k_1-1} u_i v_{i+1}$. See Fig. 1, for an 11-triangular-chain.

Let H be a $(2k - 1)$ -triangular-chain, where $k \geq 3$, and let $H_1 = H \cup \{v_k u_1, u_{k-1} v_1, v_1 v_k\}$. Then we say that H_1 is a $(2k - 1)$ -strip. See Fig. 2 for an example of an 11-strip. Note also that a 5-strip is isomorphic to K_5 . The following is easy to observe.

* Corresponding author.

E-mail addresses: roi.krakovski@cs.bgu.ac.il (R. Krakovski), cstephen@mtsu.edu (D. Christopher Stephens).




Fig. 1. An 11-triangular-chain.




Fig. 2. An 11-strip.

Lemma 1.1. Let H be a $(2k - 1)$ -strip, for some integer $k \geq 4$. Then H is claw-free, non-planar, 4-regular, 4-connected, and does not contain a subdivision of K_5 .

We can now state the main result of this paper.

1.2. Let G be a 4-connected claw-free graph. Then exactly one of the following holds:

1. G contains a subdivision of K_5 ,
2. G is planar,
3. G is isomorphic to a $(2k - 1)$ -strip, for some $k \geq 4$.

Recently, the first author has proved the following [3]. This was conjectured by Xiaoya Zha in 1992 [1].

1.3. Let G be a 4-connected non-planar line graph. Then G contains a subdivision of K_5 .

We will show that 1.3 implies 1.2.

2. Proof of main result

A k -wheel (denoted by W_k) is a graph, H , with $V(H) = \{v_0, v_1, \dots, v_k\}$ such that v_0 is adjacent to all vertices v_1, \dots, v_k , and the vertices v_1, \dots, v_k induce a cycle. The vertex v_0 is called the *hub* of the wheel.

We need the following result regarding disjoint crossing paths [4].

2.1. Let v_1, \dots, v_k be distinct vertices of a graph G . Then exactly one of the following holds.

1. there are disjoint paths of G , P_1 and P_2 , so that P_i is a p_i, q_i -path, $i = 1, 2$, and p_1, p_2, q_1, q_2 occur in the sequence v_1, \dots, v_k in order (that is, P_1 and P_2 cross with respect to v_1, \dots, v_k), or
2. there is a (≤ 3) separation (G_1, G_2) of G with $v_1, \dots, v_k \in V(G_1)$ and $|V(G_2 \setminus G_1)| \geq 2$, or
3. G can be drawn in a disk with v_1, \dots, v_k on the boundary of the disk in order.

We deduce the following.

2.2. Let G be a 4-connected non-planar graph and suppose that G contains a subgraph, $H \subseteq G$, so that one of the following holds:

1. H is isomorphic to K_4 , or
2. H is isomorphic to W_k , for some $k \geq 4$, and $\deg_G(v_0) = k$ (where v_0 is the hub of the wheel).

Then G contains a subdivision of K_5 .

Proof. We may assume that H is not isomorphic to K_4 ; for otherwise let $x \in V(G \setminus H)$ (x exists as G is non-planar). By Menger's Theorem, G has four x, H -paths, pairwise disjoint except for their end x . The union of these paths together with H is a subdivision of K_5 and the claim follows.

Hence assume that H is isomorphic to W_k , for some $k \geq 4$, and that $\deg_G(v_0) = k$. Let $V(H) = \{v_0, v_1, \dots, v_k\}$, and assume that v_1, \dots, v_k appear on the cycle $H \setminus v_0$ in order. Let $G' = G \setminus v_0$. Clearly, G' cannot be drawn in a disk with v_1, \dots, v_k appearing on the boundary of the disk in order, for since $\deg_G(v_0) = k$ (and hence all neighbors of v_0 in G are in $V(H)$) that would imply that G is planar. Also, there is no (≤ 3) -separation (G'_1, G'_2) in G' , so that $v_1, \dots, v_k \in V(G'_1)$ and $|V(G'_2 \setminus G'_1)| \geq 2$; for otherwise, all neighbors of v_0 in G are in $V(G'_1)$; but then $(G'_1 \cup v_0 \cup \{v_0 v_1, \dots, v_0 v_k\}, G'_2)$ is a (≤ 3) -separation in G , contradicting the fact that G is 4-connected.

Hence, by 2.1, in G' , there are disjoint paths P_1 and P_2 , so that P_i is a p_i, q_i -path, $i = 1, 2$, and p_1, p_2, q_1, q_2 occur in the sequence v_1, \dots, v_k in order. It is easy to see (by taking minimal crossing subpaths of P_1 and P_2) that there are subpaths $P'_1 \subseteq P_1$ and $P'_2 \subseteq P_2$ of P_1 and P_2 (possibly $P'_1 = P_1$ or $P'_2 = P_2$), such that for $i = 1, 2$, P'_i has ends $p'_i, q'_i \in \{v_1, \dots, v_k\}$, P'_i is disjoint of $\{v_1, \dots, v_k\}$ except for its ends, and p'_1, p'_2, q'_1, q'_2 appear on the sequence v_1, \dots, v_k in order. Then $H \cup P'_1 \cup P'_2 \subseteq G$ contains a subdivision of K_5 (with branch set $\{v_0, p'_1, q'_1, p'_2, q'_2\}$) and the claim follows. \square

The following result is straightforward. It follows from the fact that if G is a graph that is claw-free and triangle-free, then every vertex of G is of degree at most two.

2.3. Let G be a connected graph. Then, G is claw-free and triangle-free if and only if G is a path or a cycle of length at least four.

From 2.3 and 2.2, we obtain the following.

2.4. Let G be a 4-connected claw-free non-planar graph. If G contains a vertex of degree at least five, then G contains a subdivision of K_5 .

Proof. Let $v \in V(G)$ be a vertex of G of degree at least 5. Let H be the subgraph induced by the neighbors of v . Clearly, H is claw-free (as $H \subseteq G$ is induced). Also, we may assume that H has no triangles, for otherwise the claim follows, as then G contains a subgraph isomorphic to K_4 and hence, by 2.2, a subdivision of K_5 .

As $|V(H)| \geq 5$, H has no triangles, and $G[H \cup v] \subseteq G$ is claw-free, it must be that H is connected. To see this, assume that H is disconnected, and let C_1 be a connected component of H . Let $C_2 = H \setminus V(C_1)$. As $|V(H)| \geq 5$, there exists $1 \leq i \leq 2$, so that $|V(C_i)| \geq 3$. As H is triangle-free, there exist $a, b \in V(C_i)$ so that $ab \notin E(H)$ (and hence $ab \notin E(G)$). But then a, b and v together with any vertex of $V(C_{3-i})$ is a claw in G ; a contradiction. Hence H is connected. We also see that H is not a path; for if H is a path, say $p_1, \dots, p_{|V(H)|}$, then as $|V(H)| \geq 5$, $v, p_1, p_{\lceil |V(H)|/2 \rceil}, p_{|V(H)|}$ induce a claw in G ; a contradiction.

2.3 implies then that H is a cycle. But then $G[V(H) \cup v]$ is isomorphic to $W_{\deg_G(v)}$, and the claim follows by 2.2. \square

Proof of 1.2. Let G be a 4-connected claw-free graph. If G is planar or contains a subdivision of K_5 , then there is nothing to prove. Hence, we assume that G is non-planar and does not contain a subdivision of K_5 , and show that G is isomorphic to a k -strip, for some odd $k \geq 7$. By 2.4 and 2.2 we may assume the following.

(1) G is 4-regular and does not contain a subgraph isomorphic to K_4 .

By 1.3, G is not a line graph. Hence, since G is claw-free but not a line graph, then by [2], G contains a subgraph, H , isomorphic to a 4-triangular-chain (i.e., isomorphic to the graph obtained from K_4 by deleting a single edge). Let H be such that

- (i) H is isomorphic to a k -triangular-chain, for some $k \geq 4$;
- (ii) subject to (i), k is maximal.

Let $k_1 = \lceil k/2 \rceil$ and $k_2 = \lfloor k/2 \rfloor$. We observe that:

(2) $V(H) = V(G)$.

Subproof. We may assume that k is even (for if k is odd, the proof follows the same arguments). Let $V(H) = \{v_1, \dots, v_{k_1}, u_1, \dots, u_{k_1}\}$ with edge set as in the definition of k -triangular-chain. Note that in H , v_{k_1} is of degree three. Let $x \in N_G(v_{k_1}) \setminus \{u_{k_1}, u_{k_1-1}, v_{k_1-1}\}$. By (ii), x is not adjacent to u_{k_1} . By (1), u_{k_1} is not adjacent to v_{k_1-1} . Hence, as G is claw-free, it must be that x is adjacent to v_{k_1-1} . But this implies that $k_1 = 2$ (for then v_{k_1-1} must be of degree strictly smaller than four in H), and using x we get a contradiction to (ii).

Hence, $N_G(v_{k_1}) \subseteq V(H)$ and by the argument above we may assume that $k \geq 6$. Then, $N_G(v_{k_1}) \cap \{v_1, u_1\} \neq \emptyset$. It follows that $V(G) = V(H)$, for if there exists $v \in V(G) \setminus V(H)$, then v_1, u_1, u_{k_1} is an ℓ -cut, $2 \leq \ell \leq 3$, separating v from v_2 . This proves (2).

Hence assume that $V(G) = V(H)$. We see that k must be odd; for if k is even, then as G is 4-regular and simple it follows that $G = H \cup \{v_{k_1}v_1, u_1u_{k_1}, v_1u_{k_1}\}$, and then G is planar; a contradiction. Hence k is odd and $k \geq 5$, and we deduce as above that $G = H \cup \{v_1v_{k_1}, v_1u_{k_2}, u_2v_{k_1}\}$. Then G is isomorphic to a k -strip, for some $k \geq 5$. If $k = 5$, then G is isomorphic to K_5 ; a contradiction. Hence, $k \geq 7$, and the proof is completed. \square

References

- [1] D. Archdeacon, Problems in topological graph theory, Manuscript. Available online at: <http://www.emba.uvm.edu/archdeac/problems>.
- [2] L.W. Beineke, Characterizations of derived graphs, Journal of Combinatorial Theory 9 (2) (1970) 129–135.
- [3] R. Krakovski, Topological minors in line graphs II—Subdivisions of K_5 , Combinatorica (submitted for publication).
- [4] N. Robertson, P.D. Seymour, R. Thomas, Hadwiger's conjecture for K_6 -free graphs, Combinatorica 13 (3) (1993) 279–361.